

On the extension of 2-polynomials

Pei-Kee Lin

Department of Mathematics

University of Memphis

Memphis TN, 38152

USA

February 1, 2008

Let X be a normed linear space over $\mathbb{K}$ ($\mathbb{R}$ or $\mathbb{C}$). A function $P : X \rightarrow \mathbb{K}$ is said to be a *2-polynomial* if there is a bilinear functional $\Pi : X \times X \rightarrow \mathbb{K}$ such that $P(x) = \Pi(x, x)$ for every $x \in X$. The norm of P is defined by

$$\|P\| = \sup\{|P(x)| : \|x\| = 1\}.$$

It is known that if X is an inner product space, then every 2-polynomial defined in a linear subspace of X can be extended to X preserving the norm. On the other hand, there is a 2-polynomial P defined in a two dimensional subspace of ℓ_∞^3 such that every extension of P to ℓ_∞^3 has norm greater than $\|P\|$, (see [1, 2]). Recently, Benítez and Otero [2] showed that if X is a three dimensional real Banach space X such that the unit ball of X is an intersection of two ellipsoids, then every 2-polynomial defined in a linear subspace of X can be extended to X preserving the norm. It is natural to ask [2]

Question 1 *Suppose X is a norm space such that the unit ball of X is an intersection of two ellipsoids. Can every 2-polynomial defined in a linear subspace of X be extended to X preserving the norm?*

In this article, we show the answer is affirmative when X is a finite dimensional space.

First, we recall the following result in [2].

Lemma 1 *If P and Q are 2-polynomials in $X = \mathbb{R}^2$ such that*

$$0 \leq \sup\{P(x), Q(x)\}, \quad (x \in X)$$

then there exists $0 \leq \alpha \leq 1$ such that

$$0 \leq \alpha P(x) + (1 - \alpha)Q(x), \quad (x \in X).$$

Lemma 2 *Suppose X is a real norm space, and suppose P_1, P_2 are two positive semidefinite 2-polynomials on X . If Q is a 2-polynomial such that*

$$|Q(x)| \leq \max\{P_1(x), P_2(x)\} \quad (x \in X),$$

then there are $0 \leq \alpha, \beta \leq 1$ such that for every $x \in X$,

$$-((\beta P_1(x) + (1 - \beta)P_2(x)) \leq Q(x) \leq \alpha P_1(x) + (1 - \alpha)P_2(x).$$

Proof. For every $x \in S$ with $Q(x) \geq 0$ ($Q(x) \leq 0$), let

$$A_x = \{\gamma \in [0, 1] : \gamma P_1(x) + (1 - \gamma)P_2(x) \geq Q(x)\}$$

$$(B_x = \{\gamma \in [0, 1] : -(\gamma P_1(x) + (1 - \gamma)P_2(x)) \leq Q(x)\}).$$

To prove this lemma, it is enough to show that

$$\cap_{\{x \in S: Q(x) \geq 0\}} A_x \neq \emptyset, \text{ and } \cap_{\{x \in S: Q(x) \leq 0\}} B_x \neq \emptyset.$$

For any $x \in S$ with $Q(x) \geq 0$ (respectively, $Q(x) \leq 0$), A_x (respectively, B_x) is a nonempty closed subinterval of $[0, 1]$. So we only need to show that for any $x, y \in S$, if $\min\{Q(x), Q(y)\} \geq 0$, (respectively, $\max\{Q(x), Q(y)\} \leq 0$), then $A_x \cap A_y \neq \emptyset$, (respectively, $B_x \cap B_y \neq \emptyset$).

Suppose that $Q(x) \geq 0$ and $Q(y) \geq 0$. For any $z \in \text{span}\{x, y\}$, R_1 and R_2 are defined by

$$R_1(z) = P_1(z) - Q(z)$$

$$R_2(z) = P_2(z) - Q(z).$$

By Lemma 1, there is a γ , $0 \leq \gamma \leq 1$ such that

$$0 \leq \gamma R_1(z) + (1 - \gamma)R_2.$$

This implies $\gamma \in A_x \cap A_y$. Similarly, if $Q(x) \leq 0$ and $Q(y) \leq 0$, then there exists $0 \leq \gamma \leq 1$ such that $\gamma \in B_x \cap B_y$. We proved our lemma. $\square$

Theorem 3 *Let Π_1 and Π_2 be two inner products on $\mathbb{R}^n$ and let $X = \mathbb{R}^n$ be the space with the norm*

$$\|x\| = \sqrt{\max\{\Pi_1(x, x), \Pi_2(x, x)\}}.$$

Then every 2-polynomial defined in a subspace of X can be extended to X preserving the norm.

Proof. Let P be any 2-polynomial on a subspace Y of X and let

$$\begin{aligned}\|x\|_1 &= \sqrt{\Pi_1(x, x)} \\ \|x\|_2 &= \sqrt{\Pi_2(x, x)}.\end{aligned}$$

Without loss of generality, we may assume $\|P\| = 1$ and Y is a hyperplane of X . So for any $x \in Y$,

$$|P(x)| \leq \|x\|^2 \leq \max\{\|x\|_1^2, \|x\|_2^2\}.$$

By Lemma 2, there exist $0 \leq \alpha, \beta \leq 1$ such that for every $x \in Y$,

$$-(\beta\|x\|_1^2 + (1 - \beta)\|x\|_2^2) \leq P(x) \leq \alpha\|x\|_1^2 + (1 - \alpha)\|x\|_2^2.$$

Replacing $\|x\|_1$ (respectively, $\|x\|_2$) by $(\alpha\Pi_1(x, x) + (1 - \alpha)\Pi_2(x, x))^{1/2}$ (respectively, $(\beta\Pi_1(x, x) + (1 - \beta)\Pi_2(x, x))^{1/2}$), we may assume that if $x \in Y$, then

$$-\|x\|_2^2 \leq P(x) \leq \|x\|_1^2.$$

Let Π be the symmetric bilinear functional associated to P . Then there is a (bounded) symmetric operator T_1 (respectively, T_2) on $(Y, \|\cdot\|_1)$ (respectively, $(Y, \|\cdot\|_2)$) such that for any $y_1, y_2 \in Y$

$$\Pi(y_1, y_2) = \Pi_1(y_1, T_1(y_2)) = \Pi_2(y_1, T_2(y_2)).$$

Since T_1 and T_2 are symmetric, they are diagonalizable. It is known that

1. every eigenvalue of T_1 (respectively, T_2) is real;
2. if x_1 and x_2 are two eigenvectors of T_1 (respectively, T_2) associated with two distinct eigenvalues, then

$$\Pi(x_1, x_2) = 0.$$

Let Y_1 (respectively, Y_3) be the subspace spanned by all eigenvectors of T_1 (respectively, T_2) associated with non-negative eigenvalues, and Y_2 (respectively, Y_4) be the subspace spanned by all eigenvectors of T_1 (respectively, T_2) associated with negative eigenvalues. Then Y_1, Y_2, Y_3, Y_4 satisfy the following conditions:

3. $Y = Y_1 \oplus Y_2 = Y_3 \oplus Y_4$;
4. $T_1(Y_1) \subseteq Y_1$, $T_1(Y_2) \subseteq Y_2$, $T_2(Y_3) \subseteq Y_3$, and $T_2(Y_4) \subseteq Y_4$;
5. for any $y_i \in Y_i \setminus \{0\}$,

$$\begin{aligned}\Pi_1(y_1, T_1(y_1)) &\geq 0 > \Pi_1(y_2, T_1(y_1)), \\ \Pi_2(y_3, T_1(y_3)) &\geq 0 > \Pi_2(y_4, T_2(y_4)), \\ \Pi_1(y_1, y_2) &= 0 = \Pi_2(y_3, y_4).\end{aligned}$$

We claim that $Y_1 \cap Y_4 = \{0\}$. Suppose it is not true. Let $y \in (Y_1 \cap Y_4) \setminus \{0\}$. Then

$$0 > \Pi_2(y, T_2(y)) = \Pi(y, y) = \Pi_1(y, T_1(y)) \geq 0.$$

We get a contradiction. Similarly, $Y_2 \cap Y_3 = \{0\}$. Hence, we have

6. $\dim(Y_1) = \dim(Y_3)$, $\dim(Y_2) = \dim(Y_4)$, and

$$Y_1 \oplus Y_4 = Y = Y_2 \oplus Y_3.$$

Let

$$\begin{aligned}M_1 &= \{z \in X : \Pi_1(z, x) = 0 \text{ for all } x \in Y_1\}, \\ M_2 &= \{z \in X : \Pi_2(z, x) = 0 \text{ for all } x \in Y_4\}.\end{aligned}$$

By (5) and (6), $\dim(M_1) = \dim(Y_2) + 1$ and $\dim(M_2) = \dim(Y_3) + 1$. This implies there is a non-zero vector $z \in M_1 \cap M_2$. Let ϕ be any non-zero linear functional on X such that $\ker \phi = Y$. For any $x \in X$, define

$$\tilde{P}(x) = P\left(x - \frac{\phi(x)}{\phi(z)}z\right).$$

We claim that if $0 < \tilde{P}(x)$, then $\tilde{P}(x) \leq \|x\|^2$.

Case 1. $x - \frac{\phi(x)}{\phi(z)}z \in Y_1$. Since $z \in M_1$, we have $\Pi_1(x - \frac{\phi(x)}{\phi(z)}z, z) = 0$. So

$$\tilde{P}(x) = P\left(x - \frac{\phi(x)}{\phi(z)}z\right) \leq \left\|x - \frac{\phi(x)}{\phi(z)}z\right\|_1^2 \leq \|x\|_1^2.$$

Case 2. $x - \frac{\phi(x)}{\phi(z)}z \notin Y_1$. Then there exist $y_1 \in Y_1$ and $y_2 \in Y_2$ such that $x - \frac{\phi(x)}{\phi(z)}z = y_1 + y_2$. Note: $\Pi_1(y_1, y_2) = 0 = \Pi_1(y_1, z)$. So

$$\begin{aligned}\tilde{P}(x) &= P\left(x - \frac{\phi(x)}{\phi(z)}z\right) = \Pi(y_1 + y_2, y_1 + y_2) \\ &= P(y_1) + P(y_2) \leq P(y_1) \\ &\leq \|y_1\|_1^2 \leq \|x\|_1^2 \\ &\leq \|x\|^2.\end{aligned}$$

We proved our claim. Similarly, if $\tilde{P}(x) \leq 0$, then $|\tilde{P}(x)| \leq \|x\|^2$. Hence, $\tilde{P}$ is an extension of P preserving the norm. $\square$

In [2] (Lemma 2 and Proposition 2), Benítez and Otero proved that the problem of extension preserving the norm can be reduced to the real case. Hence, we have the following theorem.

Theorem 4 *Let Π_1 and Π_2 be two inner products on $\mathbb{C}^n$ and let $X = \mathbb{C}^n$ be the space with the norm*

$$\|x\| = \sqrt{\max\{\Pi_1(x, x), \Pi_2(x, x)\}}.$$

Then every 2-polynomial defined in a subspace of X can be extended to X preserving the norm.

Acknowledgements

The author are thankful to the referee for his comments.

References

- [1] R. M. Aron and P. D. Berner, *A Hahn-Banach extension theorem for analytic mappings*, Bull. Soc. Math. Franc. **106** (1978), 3-24.
- [2] C. Benítez and M.C. Otero, *On the extension of continuous 2-polynomials in normed linear spaces*, in “Geometric Aspects of Banach spaces”, London Math. Soc. Lecture Note Series **140**, 125-132.
- [3] K. Hoffman and R. Kunze, “Linear Algebra”, Prentice-Hall, 1971.